

Subpart E [Reserved]

Subpart F—Dermatologic Dosage Forms

§ 449.504 Amphotericin B dermatologic dosage forms.**§ 449.504a Amphotericin B ointment.**

(a) *Requirements for certification—(1) Standards of identity, strength, quality, and purity.* Amphotericin B ointment is composed of amphotericin B in a suitable and harmless ointment base. It may contain suitable and harmless coloring agents and protectants. It contains 30 milligrams of amphotericin B in each gram. Its potency is satisfactory if it is not less than 90 percent and not more than 125 percent of the number of milligrams of amphotericin B that it is represented to contain. Its moisture content is not more than 1.0 percent. The amphotericin B used conforms to the standards prescribed by § 449.4(a)(1) (i), (ii), (v), (vi), and (vii).

(2) *Labeling.* It shall be labeled in accordance with the requirements of § 432.5 of this chapter.

(3) *Requests for certification; samples.* In addition to the requirements of § 431.1 of this chapter, each such request shall contain:

(i) Results of tests and assays on:

(a) The amphotericin B used in making the batch for potency, amphotericin A content, pH, residue on ignition, and identity.

(b) The batch for potency and moisture.

(ii) Samples required:

(a) Amphotericin B used in making the batch: 10 packages, each containing not less than 500 milligrams.

(b) The batch: A minimum of 5 immediate containers.

(b) *Tests and methods of assay—(1) Potency.* Proceed as directed in § 436.105 of this chapter, preparing the sample for assay as follows: Place an accurately weighed representative portion of the sample (usually 1 gram) into an appropriate-sized Erlenmeyer flask with 10 milliliters of ethyl ether. Allow to dissolve for 1 hour with the intermittent manual shaking. Add a measured amount of dimethylsulfoxide to the flask and place on a shaker for 10 minutes. Further dilute with

dimethylsulfoxide to a concentration of 20 micrograms of amphotericin B per milliliter (estimated). Remove an aliquot and dilute with 0.2M potassium phosphate buffer, pH 10.5 (solution 10), to the reference concentration of 1.0 microgram of amphotericin B per milliliter (estimated).

(2) *Moisture.* Proceed as directed in § 436.201 of this chapter.

§ 449.504b Amphotericin B cream.

(a) *Requirements for certification—(1) Standards of identity, strength, quality, and purity.* Amphotericin B cream is composed of amphotericin B, with or without one or more suitable and harmless emollients, perfumes, dispersants, and preservatives, in a suitable and harmless cream base. It contains 30 milligrams of amphotericin B in each gram. Its potency is satisfactory if it is not less than 90 percent and not more than 125 percent of the number of milligrams of amphotericin B per gram that it is represented to contain. The amphotericin B used conforms to the standards prescribed by § 449.4(a)(1) (i), (ii), (v), (vi), and (vii).

(2) *Labeling.* It shall be labeled in accordance with the requirements of § 432.5 of this chapter.

(3) *Requests for certification; samples.* In addition to the requirements of § 431.1 of this chapter, each such request shall contain:

(i) Results of tests and assays on:

(a) The amphotericin B used in making the batch for potency, amphotericin A content, pH, residue on ignition, and identity.

(b) The batch for potency.

(ii) Samples required:

(a) Amphotericin B used in making the batch: 10 packages, each containing not less than 500 milligrams.

(b) The batch: A minimum of 5 immediate containers.

(b) *Tests and methods of assay; potency.* Proceed as directed in § 436.105 of this chapter, preparing the sample for assay as follows: With the aid of a high-speed glass blender, dissolve an accurately weighed sample in sufficient dimethylsulfoxide to give a stock solution of convenient concentration. Further dilute with dimethylsulfoxide to a concentration of 20 micrograms of